

SOV/124-57-5-5858

Translation from: Referativnyy zhurnal. Mekhanika, 1957, Nr 5, p 119 (USSR)

AUTHOR: Sokolov, N. B.

TITLE: Concerning the Stress-strain Equations Depicting Large Planar Deformations (Ob uravneniyakh svyazi mezhdur napryazheniyami i bol'shimi deformatsiyami pri ploskom deformirovannom sostoyanii)

PERIODICAL: Tr. Penzensk. industr. in-ta, 1955, Nr 3, pp 14-23

ABSTRACT: The author examines the phenomenon of planar deformation. The components of a finite deformation are expressed in terms of the displacement functions. He determines the principal elongations and the directions of the principal axes both before and after deformation. By introducing certain auxiliary quantities he is enabled to evolve an equation wherein the stress components are expressed explicitly in terms of the generalized strain components. From the expressions obtained it is possible to evolve both the Hooke equations for small elastic deformations and the Hencky equations for small elastic-plastic deformations. The stress-strain ratio can be obtained by replacing the usual strain-component term with a term adjusted for the ratio of the postdeformation surface-area increment to the

Card 1/2

SOV/124-57-5-5858

Concerning the Stress-strain Equations Depicting Large Planar Deformations

initial predeformation surface area. It is found that the tangential (shear) stress is a function of the triaxial deformation.

V. G. Osipov

Card 2/2

343

AUTHOR: Sokolov, N.B., Doctor of Technical Sciences.

TITLE: Analysis of the operation of shaft mills and generalisation of experimental data. (Analiz raboty shakhtnykh melnits i obobshenie opytnykh dannyykh.)

PERIODICAL: "Energomashinostroenie," (Power Machinery Construction), 1957, No. 3, pp. 7 - 12, (U.S.S.R.)

ABSTRACT: Evaluation and generalisation of the validity of test results obtained on industrial shaft mills during tests in crushing of coal, shale and milling peat, carried out by various authors during the last 15 to 20 years. The general preliminary analysis of the experimental data reveals the following basic features of mill operation: on feeding the fuel into the mill, operating under no-load conditions, the grinding takes place practically without any change in the required power, up to a certain limit in the quantity of the fed-in fuel. Further increase of the feeding-in of the fuel will bring about an increase in the required power, simultaneously with an increase of output; this increase in power continues up to a certain limit after which the operation of the mill becomes unstable and impossible owing to flooding of the mill by fuel. During the steady state operation of the mill, disregarding recirculation, the quantity of fed-in raw coal is equal to the quantity of pulverised coal produced by the mill. The calculation formula is based on empirical

Analysis of the operation of shaft mills and generalisation<sup>343</sup>  
of experimental data. (Cont.)

values of the experimental data, given in the graph, Fig. 5, which yields for the power consumption the formula expressed by eq. (9), p.9, and the author claims that calculations obtained by means of this formula are in good agreement with experimentally-determined values (2). The relations are also derived for the wear and service life of the crushing elements; the eq. (12), p.11, expresses the relation derived by the author for the service life. By means of this formula, nomograms can be calculated, which facilitate the selection of shaft mills for given conditions of operation and an example of such a selection has been given in an earlier paper by the author for axial mills (2). In the final part of the paper, formulae are derived for milling peat, eqs. (14) and (15), p.12, and it can be seen from eq. (15) that the specific power consumption during grinding of milling peat increases with increasing diameter and r.p.m. of the rotor, whilst, in the case of coal, the power consumption decreases with increasing r.p.m.

9 graphs, 2 tables. 6 Russian and 1 German reference.

SOKOLOV, N.B.

Relationship between the components of stress and the components of  
finite plain strain. Uch. zap. PPI no.1:17-26 '63. (MIRA 17:2)

PSHENICHNOV, Ye.A.; SOKOLOV, N.D.

Eigenvalues and the probabilities of quantum transitions in a double  
asymmetrical potential well. Opt. i spektr. 17 no.3:343-348 S '64.  
(MIRA 17:16)

KAZHUGA, V.V.; SOROLOV, N.D.

Unidimensional model of interaction between a diatomic radical  
and a solid surface. Kin. i kat. 6 no. 6:961-967 N-D '65  
(MIRA 19:1)

1. Institut khimicheskoy fiziki AN SSSR. Submitted September 9,  
1964.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200

201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300

301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400

401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500

501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600

601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700

701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800

801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900

901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100

1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200

1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300

1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400

1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500

1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600

1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700

1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800

1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900

1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000

2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100

2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200

2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212



PA 18T83

SOKOLOV, N. D.

USSR/Chemistry - Bonds, Hydrogen  
Chemistry - Reaction Rates

Jun 1946

"The Nature of the Hydrogen Bond, and the Effect of  
Complex Formation upon Reaction Rates," N. D.  
Sokolov, 2 pp

"Zhur Fiz Khim" Vol XX, No 6

Claim by the author that the reasoning shown in his  
article, spread over more sound cases, would be the  
path toward the solution of the problem of the in-  
fluence of complex formations on reaction rates.

18T83

2

**SOKOLOV, N.**

Chemical bond and characteristics of interacting systems. II. Bond energy of two atoms of different species. *N. Sokolov (Inst. Org. Chem., Moscow). Acta Physicochim. U.R.S.S. 21, 243-66 (1946) (in English); cf. C.A. 39, 13357. — Theoretical.* An extension of the previous work to the bonding between unlike atoms. The 2 parameters (effective charge of the core and effective principal quantum no.), following the method of Slater, are used to describe the bond energies of polar diatomic molecules and donor-acceptor bonding between 2 mols.

R. A. Van Nostrand

1ST AND 2ND ORDERS PROCESSES AND PROPERTIES INDEX

2

SOLOLOV, IV.

Nature of the hydrogen bond and the effect of the formation of complexes on the reaction velocity. N. Sokolov (Inst. Org. Chem., Moscow). *Acta Physico-chim. U.R.S.S.* 21, 390-2(1940)(in English).—The H bond is treated as a case of donor-acceptor binding between a polar mol. A-H and an atom contg. two paired electrons in the outer shell. The mol. ion H<sub>2</sub><sup>+</sup> is an example. R. A. Van Norderstrand

ASME-31A METALLURGICAL LITERATURE CLASSIFICATION

REGION 12345678910111213141516171819202122232425262728293031323334353637383940414243444546474849505152535455565758596061626364656667686970717273747576777879808182838485868788899091929394959697989900

C A.

2

*Nature of the hydrogen bond.* N. D. Sukelov. *Doklady Akad. Nauk S.S.S.R.* 58, 611-14(1947).—Theoretical analysis of the exchange force wave functions involved in the configuration analogous to that of the O—H...O hydrogen bond, by use of the approx. solution with the aid of Slater's functions, indicates that the H atom possesses the optimum parameters for establishment of such a link. The calcd. shift of the O—H vibration frequency is about  $320\text{ cm}^{-1}$ , which is within reasonable range of the observed  $270\text{--}490\text{ cm}^{-1}$ .  
G. M. Kosolapoff

PA 9/49T90

USSR/Physics  
Rotators  
Adsorption

Aug 48

"Natural Values and Sum State for a Semi-spherical  
Rotor," N. D. Sokolov, Moscow, 1 p

"Zhur Eksp. i Teor. Fiz." Vol XVIII, No 8

States that simultaneously with theoretical computation of adsorption coefficients and absolute speeds of heterogeneous reactions, it is necessary to calculate sum state of molecule, which is adsorbed on adsorbent surface, and at same time is able to revolve freely about the contact point. This type of

9/49T90

Aug 48

USSR/Physics (Contd)

molecule can be treated as a semi-spherical rotor.  
Submitted 29 Mar 1948.

SOKOLOV, N. D.

9/49T90

SOKOLOV, N. D.

PA 68T16

USSR/Chemistry - Hydrogen, Bonds  
Chemistry - Protons, Transfer of

May 1948

"Hydrogen Bond and Processes of Proton Transfer,"  
N. D. Sokolov, Inst Org Chem, Acad Sci USSR, 4 pp

"Dok Ak Nauk SSSR" Vol LX, No 5

Continuation of experiments first reported in "Dok  
Ak Nauk SSSR," Vol LVIII, 1947, which described  
methods for obtaining bond frequency of A....B.  
Presents functions similar to results obtained in  
first article, with which it is possible to obtain a  
satisfactory description of spectroscopic phenomena  
of hydrogen bond. Submitted by Academician A. N.  
Terenin 10 Mar 1948.

68T16

SOKOLOV, N. D.

PA 8/49T4

USSR/Chemistry - Hydrogen Ion, Discharge of Jul 48  
Chemistry - Overvoltage

"The Dependency of the Speed of the Discharge of Ions of Hydrogen on the Overvoltage," N. D. Sokolov, 3 1/2 pp

"Dok Ak Nauk SSSR" Vol LXI, No 1

As is well known, the speed of discharge of hydrogen ions on a cathode is given by the equation

$$i_{\eta} = a + b\eta \equiv a + \frac{\alpha F}{RT} \eta,$$

where  $i$  is current density,  $\eta$  is overvoltage and  $F, R, T$  have the usual meaning. The usual value of the  $\alpha$  constant is about 0.5. Sokolov deduces theoretical  
8/49T4

USSR/Chemistry - Hydrogen Ion, Discharge Jul 48  
of (Contd)

equation for  $\alpha$  and shows that it gives results in conformity with experimental value. Submitted  
14 Apr 1948.

8/49T4

CA

/ Physical bases of the theory of the chemical bond. N. D.  
Sokolov. *Uspehi Khim.* 18, 697-723(1949).—A crit.  
review, involving in particular a criticism of Pauling's  
representation of resonance. 18 references. N. Thon



Magyar Kémiai Polyoirat  
Hungarian Journal of Chemistry  
vol. 46 1950  
no. 11, november

### *N D Sociology*

On the physical foundations of the theory of chemical bonds (From the Russian) 101-120

450-524 METALLURGICAL LITERATURE CLASSIFICATION

SOKOLOV, N.D., KURSANOV, D.N., KABACHNIK, M.I., KAVERZNEVA, Ye.D., PRILEZHAYEVA, Ye.N.  
and FREYDLINA, R. Kh.

"The Current State of Chemical Structure," Usp. Khim., 19, No.5,  
pp 529-544, 1950

Translation W-16104, 30 Dec 50

BUBEN, N. YA.; VOYEVODSKIY, V. V.; SOKOLOV, N. D.

Kondrat'yev, Victor Nikolayevich, 1900-

Scientific activities of V. N. Kondrat'yev. Usp. khim. 21 no. 8, 1952.

9. Monthly List of Russian Accessions, Library of Congress, November 195~~6~~<sup>7</sup> Uncl.

SOKOLOV, N. D.

PA 227T87

USSR/Physics - Quantum Mechanics

Sep 52

"Quantum-Mechanical Treatment of the Interaction of a Polar Molecule With an Atom," N.D. Sokolov; Inst of Chem Phys, Acad Sci USSR

"Zhur Eksper i Teoret Fiz" Vol 23, No 3, pp 315-326

On the basis of the method of electron pairs, the author obtains an approx soln for the problem of interaction of a polar di-atomic mol with an atom possessing a filled external shell. The results are applied to the treatment of the hydrogen bond. Received 1 Aug 51.

227T87

SOKOLOV, N.D.

U.S.S.R.

Theory of spectroscopic phenomena of the hydrogen bond. N. D. Sokolov. *Zhur. Eksp. i Teor. Fiz.* 23, 592-603(1952); *Science Abstr.* 56A, 726(1953); cf. *C.A.* 46, 6070f. — Among the unavoidable factors which det. the magnitude of the frequency shift is not only the dependence of  $\nu$  on  $R_{AH}$ , but also the lengthening of the bond A—H during formation of the system. This extension requires anharmonicity of vibrations. K. L. C.

Sokolov, N.D.

USSR.

Quantum mechanical determination of the vibration frequencies of a compound with a hydrogen bond. N. D. Sokolov. *Zhur. Eksptl. i Teoret. Fiz.* 23, 404-11 (1953); *Soviet Phys. JETP* 17, 278-84 (1953).—The vibration frequencies of the linear triat. system A—H...B with a H-bond were investigated. The most probable optical transition corresponds to a frequency differing little (~20%) from that obtained classically. An approx. proportionality exists between the A—H frequency shift under compd. formation and the energy of the H-bond. K. L. C.

Chem 309

RM  
RM

SOKOLOV, N. D.

Quantum mechanical calculations of the deviations from the additivity scheme of organic chemistry. N. D. Sokolov (Inst. Chem. Phys., Acad. Sci. U.S.S.R., Moscow). *Dokl. Akad. Nauk SSSR*, 1952, 163, 1036-9 (1952).—The additivity scheme of org. chemistry (cf. Syrkin, *C.A.* 36, 393\*) which has been developed for satd. and ethylene hydrocarbons does not hold for aromatic compds. or those having a conjugated bond. In the latter cases the true heats of formation are usually higher than are those calcd. by the additive scheme. The deviation from this system has been calcd. by the methods of quantum mechanics. Only the  $\pi$  electrons are taken into account in the calcns., and the calcns. are carried out with a precision of some empirical factor. This factor appears to be almost const. for different mols. This is true for both the method of electron pairs and the method of mol. orbitals. J. Rovtar Leach

Quantum theory of the spectroscopic manifestations of the hydrogen bond. N. D. Sokolov (Inst. Chem. Phys. Acad. Sci. U.S.S.R., Moscow). *Doklady Acad. Nauk S.S.S.R.* 82, 372-72 (1952).—For the H bond  $RA \cdots HB$  (where A, B = N, O, F), if the interaction energy  $w$  of the group AH with the atom H is written in Jacobi coordinates  $w = w(R - ar)$ , with the distances  $r = R_{AH}$  and  $R = aR_{AH} + R_{BH}$ , and  $a = m_B/(m_A + m_B)$ , and it is assumed that  $w$  depends only on  $R_{AH}$  and not directly on  $r$ , it is found that the vibration frequency of the A—H group is shifted to shorter waves, which is in conflict with the observed fact of a shift to longer waves by a few hundreds of  $\text{cm}^{-1}$ . If a direct electrostatic interaction of the effective charges of A and B is taken into account, the frequency shift is practically zero. A shift consistent with the stationary states of the system. Numerical estimation shows the relative frequency shift to be proportional to the energy  $\epsilon$  of the H bond, with a proportionality factor of  $\sim 1.0 \times 10^{-4}$  and  $1.2 \times 10^{-5}$ , estd. from exptl. data, resp., by Shubin (C.A. 45, 3245) and Badger and Bauer (C.A. 32, 167). If, instead of taking into account a variation of the effective charge of the H atom with  $r$ , of the form  $w \sim r^{-\beta}$  (where  $\beta$  = fraction of electron assigned to the H atom), one assumes it to be constant,  $w$  becomes an ordinary Morse function of  $R_{AH}$ , and the sign of the frequency shift is in contradiction with the facts. The conclusion of these considerations can be expressed in the form that, with increasing distance A—H, the H atom undergoes some extent of protonization. N. Thon



SOKOLOV, N. D.

Defended his Dissertation for Doctor of Physicomathematical Sciences, Leningrad State University, Leningrad, 1953.

Dissertation: "Intermolecular Reactions and Processes of Proton Transition"

SO: Referativnyi Zhurnal Khimii, No. 1, Oct. 1953 (W/29955, 26 Apr 54)

SOKOLOV N.D.

TERENIN, A.N., akademik; KONDRAT'YEV, V.N., akademik; KNUNYANTS, I.L., akademik; KABACHNIK, M.I.; SOKOLOV, N.D., doktor fiz.-mat. nauk; REUTOV, O.A., doktor khimicheskikh nauk; MOSKVICHEVA, N.I., tekhnicheskiiy redaktor

[Status of the theory of chemical structure in organic chemistry]  
Sostoianie teorii khimicheskogo stroeniia v organicheskoi khimii.  
Moskva, Izd-vo Akademii nauk SSSR, 1954. 122 p. [Microfilm]

(MLRA 7:10)

1. Chlen-korrespondent AN SSSR (for Kabachnik) 2. Akademiya nauk  
SSSR. Otdeleniye khimicheskikh nauk  
(Chemical structure) (Chemistry, Organic)

VOL'KENSHTEYN, M.V.; SOKOLOV, N.D., professor, redaktor; BSHMAN, Yu.A.  
redaktor; SMIRNOVA, A.V., tekhnicheskii redaktor.

[Molecules and their structure] Molekuly i ikh stroenie. Moskva,  
Izd-vo Akademii nauk SSSR, 1955. 229 p. (MLRA 8:12)  
(Molecules)

AF701597

TREASURE ISLAND BOOK REVIEW

AID 809 - S

SOKOLOV, N. D., (Institute of Chemical Physics)

O VOZMOZHNOY ROLI DONORNO-AKTSEPTORNOY SVYAZI V GETEROGENNOM KATALIZE

(The potential role of the donor-acceptor bond in heterogeneous catalysis).

In Problemy kinetiki i kataliza (Problems of Kinetics and Catalysis), vol. 8.

Izdatel'stvo Akademii Nauk SSSR, 1955. Section II: General problems of the theory of catalysis. p. 141-144.

The adsorption of molecules by solids might be explained by a donor-acceptor bond, i.e., as a result of the "polarization" of the molecule-"donor" in a positive -ion field or an atom (with a positive charge) of a molecule (crystal). However, there are few experimental data confirming the adsorption of molecules by means of donor-acceptor interaction. (A. N. Terenin's work on spectra of ammonia, some amines and other substances adsorbed by heteropolar crystals). There are several papers dealing with the theory of catalysis in which it is assumed that donor-acceptor bonds are formed in various stages of chemical transformations. The quantum - mechanism treatment of the donor-acceptor bond leads to the conclusion that the energy of the bond depends on the parameters which characterize the state of the non-shared electrons. The elucidation of the donor-acceptor mechanism should be confirmed experimentally, namely by studying spectra of adsorbed molecules and the poisoning of catalysts by impurities known to be absorbed with the aid of the donor-acceptor bond. 9 refs., 6 Russian (1929-1953).

1/1

SOKOLOV, N.D.

Possible role of the donor-acceptor bond in heterogeneous catalysis.  
Probl.kin.i kat. 8:141-144 '55. (MLRA 9:5)

1. Institut khimicheskoy fiziki AN SSSR.  
(Catalysis)

*K* *N.D.*  
*S. Am...*  
The hydration heat of atoms. V. N. Kondrat'ev and N. D. Sokolov (Inst. Chem. Phys., Acad. Sci. U.S.S.R., Moscow). Zhur. Fiz. Khim. 29, 1200-72 (1955).—A system of relative heats of ion hydration is proposed, based on the universally accepted system of heat of formation of ions in soln., in which the heat of formation of the H ion in a dil. soln. is equated to 0. The heats of hydration of some alk-oxy ions were calcd. on the basis of the relation between the hydration heats of the ions and their radii; also the affinity of the simplest alc. mol., to protons, etc. W. M. S.

*2*  
*Sum* *Est*

Sokolov, N.D.

2

The hydrogen bond. N. D. Sokolov. *Uspekhi Fiz. Nauk* 57, 205-78(1955).—A review covering the exptl. and theoretical work on the H bond including its appearance in vibration, electron, and magnetic spectra. 130 references. J. Rovtar Leach

CH  
MA  
WST

SOKOLOV, N.D.

Hydrogen bond in proton transfers. Ukr. khim.zhur. 22 no.1:18-22  
'56. (Hydrogen) (Protons) (MLRA 9:6)



SOKOLOV, N.D.

2294. CORRELATION OF NUCLEAR QUADRUPOLE INTER- 539.132  
ACTION FREQUENCIES WITH POLAROGRAPHIC HALF-  
WAVE POTENTIALS. T. Jredale.  
Nature (London), Vol. 117, 36-7 (Jan. 7, 1956).

2295. THE HYDROGEN BOND. N.D. Sokolov. 539.132  
Uspekhi fiz. Nauk, Vol. 57, No. 2, 264-76 (1957). In  
Russian.

A survey article by the leading Soviet worker in the field.  
Among topics discussed are the following:- vibration spectra,  
the inadequacy of the simple electrostatic conception of the H-  
bond, quantum-mechanical treatment, calculation of the vibrat-  
ion frequency of an AH group in a complex with a hydrogen  
bond, the origin of the band in the vibration spectrum of an H-  
bond, electric and magnetic spectra, H-bond and proton tran-  
sition processes. 139 refs. C.R.S. Manders

*Sokolov, N. D.*  
7  
3  
1-453  
Interaction between hydroxyl groups and aromatic com-  
pounds. N. D. Sokolov (Inst. Phys. Chem., Acad. Sci.  
U.S.S.R., Moscow). *Priroda*, Khim. Soderzh. i Reaktsii  
Yadrenykh Adsorbentov, Sbornik Trudov Konferentsii Adsorbentov  
1957, 257-9. - A discussion. A. P. Kotloby

3050207 A.D.

51-5-26/26

AUTHOR: Sokolov, N.D.

TITLE: Letter to the Editor: On the Article of V.M.Chulanovskiy and M.P.Burgova "On Inter-molecular Interaction in Solutions of Chloroform and Bromoform", II. (Ref.1)  
(Pis'mo v redaktsiyu. Po povodu stat'i V.M.Chulanovskogo i M.P. Burgovoy "O mezh-molekulyarnom vzaimodeystvii v rastvorakh khloroforma i bromoforma. II")

PERIODICAL: Optika i Spektroskopiya, 1957, Vol.2, Nr 5, p.680 (USSR)

ABSTRACT: This is a complete translation. In using an arbitrary model of a hydrogen bond  $A-B...C$  ( $B = \text{hydrogen atom}$ ) it is necessary for the energy of the system to be minimum at  $r = r'_0$  and  $R = R'_0$ . If this condition is not satisfied then one can obtain any result one wants to. From the basic equation in the article of V.M.Chulanovskiy and M.P. Burgova (Ref.1)  $F(r, R-r) = F(r) - f(r) - \psi(R-r)$  it follows that this condition is not obeyed, since  $\psi(R-r)$  includes only force of attraction between the atoms B and C but it does not contain the mutual repulsion of the latter and neglects the interaction between A and C altogether. For these reasons, therefore, the system  $A-B...C$  is unstable. Thus the authors' conclusions leading in particular to the correct dependence of the frequency displacement ( $\Delta\nu$ ) on

Card 1/3

51-5-26/26

Letter to the Editor: On the Article of V.M.Chulanovskiy and M.P.Burgova "On Inter-molecular Interaction in Solutions of Chloroform and Bromoform.II".

the increase of the equilibrium length of the bond A-B (Ref. 2,3) do not prove anything. It is shown (Ref.2,3) that if the condition of equilibrium is taken into account in the expression for the energy [i.e., in  $\varphi(R-r)$  the repulsive force between B and C is included] then the frequency displacement is the reverse of that obtained experimentally. It is also shown that when the direct interaction of A and C is taken into account, then the value of  $\Delta\nu$  is obtained with a correct sign but it is one order smaller than the experimental value. These results are not paradoxical as the authors of Ref.1 think, but they only mean (Ref.2,3) that an arbitrary model of a hydrogen bond which leads to expressions for the energy similar to, say, those given by an electrostatic model, is insufficient to explain the frequency displacement of the group AB (OH) in formation of this bond. Successful results are only possible by using quantum mechanics (Ref.2 to 4) which leads to more complicated dependence of the energy on interatomic distances in the system A-B...C and makes it possible to find the factors which cause the frequency displacement. The calculations show that the

Card 2/3

SOKOLOV, N. D.

VAYNSHTEYN, L.A., kandidat fiziko-matematicheskikh nauk; SOKOLOV, N.D.,  
doktor fiziko-matematicheskikh nauk.

Theoretical spectroscopy and quantum mechanics of molecules.  
(Conference in Moscow). Vest. AN SSSR 27 no.6:101-104 Je '57.  
(MIRA 10:7)  
(Quantum theory--Congresses) (Spectrum, Molecular)

GOROTOV, N.D.,

"On the Quantum Theory of the Hydrogen Bond," paper submitted at IUPAP Symposium on Nature of Hydrogen Bonding, Ljubljana, Yugoslavakis, 30 July - 3 Aug 57.

Trans. Encl. B-3,096,177, 20 Jan 58.

also in Vestnik AN SSSR, 1957, Vol. 27, Nr. 11, pp. 137-139, "An Intl. Symposium on the Hydrogen Bond in Ljubljana." by Vol'kenshteyn, M. V.

SOKOLOV, N.D.

Relation between activation energy and the heat of reaction.

Dokl. AN SSSR 112 no.4:710-713 F '57. (MLRA 10:4)

1. Institut khimicheskoy fiziki Akademii nauk SSSR. Predstavleno akademikom V. N. Kondrat'yevym.

(Chemical reaction, Heat of)

(Activity coefficients)

Сокотов, Н. Д.

KURAREV, S. I. and SOKOLOV, N. D.

"On the Theory of Dependence of Molecular Spectra on Intermolecular Interaction."

report presented at the 4th International Meeting of Molecular Spectroscopy, Bologna,  
Italy, 7-12 Sept 1959.

Moscow State University, Moscow, USSR.



21(1), 24(7)

SOV/51-6-5-25/34

AUTHORS: Glasko, V.B., Maslov, V.P., Panikar, V.I. and Sokolov, N.D.

TITLE: On the Type of Correlation Function for the Helium Atom (О виде корреляционной функции для атома гелия)

PERIODICAL: Optika i Spektroskopiya, 1959, Vol 6, Nr 5, pp 698-700 (USSR)

ABSTRACT: In molecular calculations correlation in the motion of electrons is allowed for by introducing into the wave-function an additional factor dependent on inter-electron distance  $r_{ij}$  (Ref. 1). In analogy with the first approximation in the helium atom calculations carried out by Hylleraas (Ref 2), this multiplier can be written for a two-electron system in the form

$$f(r_{12}) = 1 + d r_{12} \quad (1)$$

where  $d$  is a variational parameter. In the general case the correlation function should depend on three correlation variables and  $f$  can be then represented as a series in powers of these variables (Refs 2, 3). When only one correlation variable is used the choice of the function  $f(r_{12})$  in the form given by Eq (1) is an arbitrary one. The question arises as to whether this choice is the best possible one. This question is answered by determining the correlation function  $f(r_{12})$  for the helium

Card 1/2

SON/51-6-5-25/34

On the Type of Correlation Function for the Helium Atom

atom by a variational method. The result is shown as curve I in a figure on p 700; curve II represents the Hylleraas function given by Eq (1). Both curves are plotted as functions of distance in atomic units. The figure shows clearly that the correlation function approximation in the form of Eq (1) is practically the best choice, at least for atoms. The paper is entirely theoretical. There are 1 figure and 6 references, 3 of which are Soviet, 1 English, 1 German and 1 mixed (Soviet, English and French).

SUBMITTED: November 29, 1958

Card 2/2

SOV/76-33-3-4/41

5(4)

AUTHOR:

Sokolov, N. D.

TITLE:

On the Energy of Conjugation (Ob energii sopryazheniya)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 3,  
pp 534 - 542 (USSR)

ABSTRACT:

It is proved that a molecule model in which only the  $\pi$ -electrons are taken into account for the purpose of clarifying the decrease in energy in connection with the formation of conjugate bonds is insufficient as according to this model an increase in energy would have to take place instead of a decrease in energy. This is the reason why in connection with the aforesaid effect it is necessary to consider also the  $\sigma$ -electrons apart from the  $\pi$ -electrons. The energy of interaction between two  $\sigma$ -bonds or one  $\sigma$ -bond and one  $\pi$ -bond - which are usually neglected in connection with calculations of the deviations from the additive scheme - are of the same order of magnitude as the energy of the interaction between two  $\pi$ -bonds. As one of the main factors causing a decrease in energy in connection with the formation of con-

Card 1/2

## On the Energy of Conjugation

SOV/76-33-3-4/41

jugate bonds the fact is regarded that the energy of the covalent bond decreases in the case of interaction with the other bond or with the free electron of the adjacent atom. According to a simplified variant of the method of the valence structure (using the equation according to London (1)) the above effects are qualitatively worked out by means of the example of the transition from propane to propylene. In connection with the aforesaid it is possible to explain the decrease in energy which is used in the case of breaking of the C-H bond in the  $\text{CH}_3$ -group

and which decreases in connection with the transition from propane to propylene to 18 kcal; i.e. the energy of interaction between the free electron and the  $\pi$ -bond C=C in allyl is smaller than in the case of the  $\sigma$ -bond C-H in propyl, as the  $\pi$ -bond is weaker than the  $\sigma$ -bond. There are 2 figures and 14 references, 8 of which are Soviet.

ASSOCIATION: Akademiya nauk SSSR, Institut khimicheskoy fiziki Moskva (Academy of Sciences, USSR, Institute of Chemical Physics Moscow)  
SUBMITTED: August 16, 1956  
Card 2/2

5 (4)

AUTHORS:

Vetchinkin, S. I., Pshenichnov, Ye. A., SOV/76-33-6-16/44  
Sokolov, N. D.

TITLE:

Influence of the Hydrogen Bond on the Energy of the Ion  
Lattice of Ammonium Chloride and Evaluation of the Affinity  
of Ammonia Molecules to the Proton (Vliyaniye vodorodnoy  
svyazi na energiyu ionnoy reshetki khloristogo ammoniya i  
otsenka srodstva molekuly ammiaka k protonu)

PERIODICAL:

Zhurnal fizicheskoy khimii, 1959, Vol 33, Nr 6,  
pp 1269-1274 (USSR)

ABSTRACT:

It may be assumed that in ion crystals containing  $H_3O^+$  or  $NH_4^+$   
ions, between cation and anion beside the Coulomb forces there  
is a hydrogen bond which increases the stability of the ion  
lattice. Usually, in energy computations this hydrogen bond  
is not considered (e.g. reference 1), which leads to a lesser  
result in computations of ion lattice energy. If, however, the  
exact ion lattice energy (IE) is known, the important  
molecular constant - the proton affinity (P) of the molecule -  
may be computed according to equation (1). The recently  
obtained value of Ref 3 for the (P) of the water molecule

Card 1/3

Influence of the Hydrogen Bond on the Energy of the Ion Lattice of Ammonium Chloride and Evaluation of the Affinity of Ammonia Molecules to the Proton SOV/76.33-6-16/44

is lower by 19 kcal as compared to the value obtained according to Ref 2, which points to the fact that in the computations per (Ref 3) the effect of the hydrogen bond between cation and anion was neglected. From quantum-mechanical computations (Ref 5) of the energy of interaction of the molecule A - H with the atom B (which exhibits an undivided electron pair) the following equation was derived:  $W = Q + P_1\omega - P_2$  (2)

( $Q$  = Coulomb energy,  $P_1\omega$  = repulsive energy between H and B,  $P_2$  = exchange (or donor-acceptor) energy of the attraction

between H and B). An investigation is then made of the applicability of equation (2) to the computation of

interaction between the cation  $NH_4^+$  and anion  $Cl^-$  in the  $NH_4Cl$  crystal and it was found that by the selected semiempirical computation method a computation is possible only if  $P_2 = 0$

is assumed, by which a lower (IE) is obtained. The change of the (IE) caused by the hydrogen bond is assumed to be of the

Card 2/3

Influence of the Hydrogen Bond on the Energy of the Ion Lattice of Ammonium Chloride and Evaluation of the Affinity of Ammonia Molecules to the Proton SOV/76-33-6-16/44

same magnitude as the last mentioned decrease in the (IE). From this point of view a computation of the lattice energy for ammonium chloride is made and it is found that the correction of the computation according to Bleick (Ref 1), in which the hydrogen bond was neglected, must be of the magnitude 10 kcal, and, consequently, the value  $P_{NH_3} = 194 \pm 7$  kcal. There are 1 figure, 1 table, and 10 references, 7 of which are Soviet.

ASSOCIATION: Akademiya nauk SSSR, Institut khimicheskoy fiziki, Moskva (Academy of Sciences of the USSR, Institute of Chemical Physics, Moscow)

SUBMITTED: October 31, 1957

Card 3/3

5(4)

AUTHORS:

Aleksandrov, I. V., Sokolov, N. D.

SOV/20-124-1-32/69

TITLE:

The Hydrogen Bond and Proton Magnetic Resonance (Vodorodnaya svyaz' i protomnyy magnitnyy rezonans)

PERIODICAL:

Doklady Akademii nauk SSSR, 1959, Vol 124, Nr 1, pp 115-118 (USSR)

ABSTRACT:

The dependence of the shielding of a proton upon the degree of the polarity of the A-H-bond can be estimated according to a method suggested by I. V. Aleksandrov (Refs 10, 11). If, as a wave function, the molecular path composed of the 1s-function of the H-atom as well as of the 2s- and 2p-Slater functions of the O-atom is selected in the form

$$\psi^0 = N [\psi_{1s} + \lambda (a \psi_{2s}^0 + b \psi_{2p}^0)], \text{ the parameter } \lambda$$

characterizes the degree of polarity of the bond O - H. The value of  $\lambda$  is near 1 and the constant  $\sigma$  of magnetic shielding can be determined as a function of  $\lambda$ . At the distance  $R = 1 \text{ \AA}$  between the nuclei it holds with sufficient accuracy that

$$\Delta \sigma_1 = -1.5 \cdot 10^{-5} \Delta \lambda. \text{ If } \lambda \text{ increases by } \Delta \lambda = 0.3 \text{ under the influence of the H-bond, it holds that } \Delta \sigma_1 \approx -4.5 \cdot 10^{-6}. \text{ The}$$

Card 1/3



The Hydrogen Bond and Proton Magnetic Resonance

SOV/20-124-1-32/69

influence exercised by expansion of the O-H-bond can be estimated by means of the same formulas as the influence exercised by polarity. For small variations of R in the case of  $\lambda = 1$  it holds that  $\Delta\sigma'_2 = -k \cdot 10^{-5} \Delta R$ . At  $\Delta R = 0.05 \text{ \AA}$  one

finds  $\Delta\sigma'_2 \sim -0.5 \cdot 10^{-6}$ . The influence of the donor-acceptor bond  $H^+ \dots \dots \dots O$  can be approximately taken into account if the molecular orbit of this bond is represented in the form

$\psi = N(\psi_{2p} + \beta\psi_{1s})$ . If the inter-atomic distance  $H \dots \dots O$  is  $1.7 \text{ \AA}$ , the contribution of this bond made towards magnetic shielding of the proton amounts to

$\Delta\sigma'_3 = (-0.08 + \beta + 3.6\beta^2) \cdot 10^{-5}$ , and with  $\beta \approx 0.2$  there follows

$\Delta\sigma'_3 \approx +2.5 \cdot 10^{-6}$ . If an electric field is applied  $\sigma$  varies by

$\Delta\sigma'_4 = -(5a_o^3/mc^2) E^2$ , where  $a_o$  denotes the Bohr radius. The

influence exercised by the second not filled electron pair (the cloud of which has an axis which is vertical to the line  $H \dots \dots O$ ) of the O-atom is taken into account by the formula

Card 2/3

The Hydrogen Bond and Proton Magnetic Resonance

SOV/20-124-1-32/69

$\Delta\sigma_5 = \frac{\chi - \chi_{zz}}{R^3}$ , where  $\chi_{zz}$  denotes the  $zz$ -component of the diamagnetic susceptibility  $\chi$ . In the case of  $R = 1.7 \text{ \AA}$ ,  $\Delta\sigma_5$  is in the interval of from  $+0.2 \cdot 10^{-6}$  to  $-0.2 \cdot 10^{-6}$ .

The data concerning the magnetic shielding of the proton in the H-bond confirm, firstly, the hypothesis that the polarity of the A-H-bond increases considerably by the formation of an H-bridge and, secondly, they agree with the hypothesis of the formation of the donor-acceptor-bond AH...B. There are 11 references, 3 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR ( Institute for Chemical Physics of the Academy of Sciences, USSR)

PRESENTED: July 28, 1958, by V. N. Kondrat'yev, Academician

SUBMITTED: July 23, 1958

Card 3/3

5(4), 24(7)

SOV/20-124-2-35/71

AUTHORS: Nikitin, Ye. Ye., Sokolov, N. D.

TITLE: On the Relation Between the Dissociation Constants of the Thermal Decomposition of Biatomic Molecules in the Presence and in the Absence of Equilibrium (O sootnoshenii mezhdu konstantami skorosti termicheskogo raspada dvukhatomnykh molekul pri nalichii i pri otsutstvii ravnovesiya)

PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 124, Nr 2, pp 366-369 (USSR)

ABSTRACT: Short reference is first made to earlier papers on the dissociation of biatomic molecules  $AB + M \rightarrow A + B + M$ . The disturbance of Boltzmann distribution over the oscillation states of the decomposing molecule decreases the degree of population of the oscillation levels near the limit of dissociation. The disturbance of Boltzmann distribution increases as a result of the decrease of the oscillation quanta near the limit of dissociation. The kinetics of thermal decomposition and of recombination probably does not satisfy the usual kinetic equation. Dissociation is a process developing in several stages and therefore the kinetics of decomposition and recombination

Card 1/4

SOV/20-124-2-35/71

On the Relation Between the Dissociation Constants of the Thermal Decomposition of Biatomic Molecules in the Presence and in the Absence of Equilibrium

must be described by the following systems of equations:

$$(d/dt)x_n = - \sum_k P_{nk} x_n + \sum_k P_{kn} x_k. \text{ Here } x_n \text{ denotes the distribu-}$$

tion function, i.e. the degree of population of the n-th level of the system AB, and  $P_{kn}$  - the probabilities of the rotation- and oscillation transitions  $k \rightarrow n$  of the molecule AB. The sums of the formula given above extend also to states of the continuous part of the spectrum. The authors investigate the simplest case in which at  $[AB] \ll [M]$  the concentrations  $[A]_0$  and  $[B]_0$  are kept constant. The aforementioned system of equations becomes linear and can be written down in the form

$(d/dt)\hat{X} = \hat{B}\hat{X} + \hat{C} [A]_0 [B]_0$ . The only column of the matrix  $\hat{X}$  is composed of the components of the distribution function  $x_n$ , and the matrices  $B$  and  $C$  describe the successive decomposition and recombination respectively. An expression is then written down for the general solution of the last-mentioned equation.

Card 2/4

SOV/20-124-2-35/71

On the Relation Between the Dissociation Constants of the Thermal Decomposition of Biatomic Molecules in the Presence and in the Absence of Equilibrium

The first eigenvalue of the matrix  $-\hat{B}$  corresponds to the dissociation-constant of decomposition and is, in the case of lacking recombination, determined by the expression  $\mu_0 = -(d/dt) [AB] / [AB]$ . The second eigenvalue  $\mu_1$  of the matrix  $-\hat{B}$  is approximately equal to the rate of relaxation of the oscillation energy of the molecule AB, where, by the way,  $\mu_0 \ll \mu_1$  holds. The solution of the equation  $\hat{B}\hat{X}_0 + \hat{C}[A]_0[B]_0 = 0$  is the distribution function  $x_n^{(0)} = N \exp(-E_n/kT)$  corresponding to equilibrium. Next, the amount of the constant N is determined, after which an equation is written down for the kinetics of thermal decomposition and of recombination. In the case of equilibrium the dissociation constant of decomposition is greater than the dissociation constant  $k_d$  in the case of lacking equilibrium. The calculated values of  $k_d$  for the thermal decomposition of  $J_2$ ,  $Br_2$  and  $O_2$  agree satisfactorily with the

Card 3/4

SOV/20-124-2-35/71

On the Relation Between the Dissociation Constants of the Thermal Decomposition of Biatomic Molecules in the Presence and in the Absence of Equilibrium

values which were determined experimentally by the kinetic method in the decomposition of molecules in the shock wave in an argon atmosphere at temperatures of 1500 - 4000° K. The dissociation constant corresponding to equilibrium is greater by one order of magnitude than the value corresponding to the lacking equilibrium. There are 15 references, 5 of which are Soviet.

ASSOCIATION: Institut khimicheskoy fiziki Akademii nauk SSSR (Institute for Chemical Physics of the Academy of Sciences, USSR)

PRESENTED: August 7, 1958, by V. N. Kondrat'yev, Academician

SUBMITTED: July 11, 1958

Card 4/4

S/051/60/008/04/028/032  
E201/E691

AUTHORS: Aleksandrov, I.V., Korst, N.N. and Sokolov, N.D.

TITLE: The Effect of the Mobility of Protons on the Width of Nuclear Magnetic Resonance Lines in Crystals

PERIODICAL: Optika i spektroskopiya, 1960, Vol 8, Nr 4, pp 575-577 (USSR)

ABSTRACT: The mobility of hydrogen atoms (protons) in condensed phases may be due to internal rotation or translational transitions from one equilibrium position to another. Nuclear magnetic resonance of protons is one of the most effective methods of investigation of their mobility. The present note describes how the second moment of the nuclear magnetic resonance signal of protons in ice can be used to find the mechanism of proton transitions. The paper is entirely theoretical. There are 6 references, 5 of which are English and 1 from Acta Crystallographica.

SUBMITTED: October 10, 1959

Card 1/1

84895

S/048/6C/024/008/018/018/XX  
B013/B067

Theory of the Thermal Decay of Diatomic  
Molecules

than  $D$ . By integrating (8), an additional pre-exponential factor appears. The non-equilibrium distribution function corresponding to the exponential decay is determined by the first eigenvector  $l_n(\mu_0)$  of the matrix  $B$ .  $l_n(\mu_0)$  can be easily found from the general formulas of the eigenvectors of the Jacobian matrix if the formula for  $\mu_0$  is known. The percentual improvement in population of the upper vibration levels is of the order of  $\exp[(D - E_p)/kT]$ , where  $p$  is the number of vibration levels calculated from the dissociation limit. This result especially corresponds to Refs. 9 and 13 dealing with the model of a decaying harmonic oscillator. Since in the case of a real anharmonic oscillator the oscillation quanta at the dissociation limit are small compared with  $kT$ , numerous levels reach the region of great impoverishment  $D - E_p \approx kT$ . Hence, it may be said that the equilibrium of the Boltzmann distribution in decay reactions is always perturbed. Since dissociation is a complex multistage process, the quantity  $K_d$  becomes small compared with the quantity obtained from the equilibrium condition  $(K_d)_{\text{equil}} = K K_{\text{rec}}$  owing to the perturbation of the

Card 2/3



S/058/61/000/011/005/025  
A058/A101

AUTHORS: Aleksandrov, I.V., Korst, N.N., Sokolov, N.D.

TITLE: Proton exchange effect on nuclear magnetic resonance line width in crystals

PERIODICAL: Referativnyy zhurnal. Fizika, no. 11, 1961, 126, abstract 11V221 (V sb. "Paramagnitn. rezonans", Kazan', Kazansk. un-t, 1960, 186 - 188)

TEXT: The authors calculate the second moment of the nuclear magnetic resonance line in ice crystals. They show that incident to proton tunneling along the hydrogen bonds the second moment decreases by  $\sim 20\%$  (on condition that the tunneling frequency is appreciably greater than the nuclear magnetic resonance line width).

I. Aleksandrov

[Abstracter's note: Complete translation]

Card 1/1

PSHENICHNOV, Ye.A.; SOKOLOV, N.D.

Quantum effects in a double potential well and spectroscopy  
of the hydrogen bond. Opt.i spektr. 11 no.1:16-23 J1 '61.  
(MIRA 14:10)  
(Hydrogen bonding) (Quantum theory) (Spectrum analysis)

VETCHINKIN, S.I.; SOKOLOV, N.D.

Calculation of the system  $\text{HeH}^+$  by means of the valence structure method. Zhur. fiz. khim. 35 no.7:1645-1647 J1 '61.

(MIRA 14:7)

1. AN SSSR, Institut khimicheskoy fiziki.  
(Helium hydride)

PSHENICHNOV, Ye.A.; SOKOLOV, N.D.

Polanyi's law for proton transitions and the hydrogen bond. Dokl.  
AN SSSR. 137 no. 2:352-355 Mr '61. (MIRA 14:2)

1. Institut khimicheskoy fiziki AN SSSR i Moskovskiy gosudarstvennyy  
universitet im.M.V. Lomonoseva. Predstavleno akademikom V.N.  
Kondrat'yevym.

(Protons) (Hydrogen bonding)

FRISH, S.E., otv. red.; BOBOVICH, Ya.S., kand. fiz.-matem. nauk, red.;  
VOL'KENSHTEYN, M.V., doktor fiz.-matem. nauk, red.; GALANIN,  
M.D., doktor fiz.-matem. nauk, red.; DRUKAREV, G.F., doktor  
fiz.-matem. nauk, red.; YEL'YASHEVICH, M.A., akademik, red.;  
KALITEYEVSKIY, N.I., doktor fiz.-matem. nauk, red.; KUSAKOV,  
M.M., doktor khim. nauk, red.; LIPIS, L.V., doktor tekhn.nauk,  
red.; PEKAR, S.I., doktor fiz.-matem. nauk, red.; PROKOF'YEV,  
V.K., doktor fiz.-matem. nauk, red.; SOKOLOV, N.D., doktor  
fiz.-matem. nauk, red.; FEOFILOV, P.P., doktor fiz.-matem.  
nauk, red.; CHULANOVSKIY, V.M., doktor fiz.-matem. nauk, red.;  
SHPOL'SKIY, E.V., doktor fiz.-matem. nauk, red.; YAROSLAVSKIY,  
N.G., kand. fiz.-matem. nauk, red.; LEKSINA, I.Ye., red. izd-  
va; PENKINA, N.V., red. izd-va; NOVICHKOVA, N.D., tekhn. red.;  
KASHINA, P.S., tekhn. red.

[Physical problems in spectroscopy] Fizicheskie problemy spektro-  
skopii; materialy. Moskva, Izd-vo Akad. nauk SSSR. Vol.1. 1962.  
474 p. (MIRA 16:2)

1. Soveshchaniye po spektroskopii. 13th, Leningrad, 1960. 2. Chlen-  
korrespondent Akademii nauk SSSR (for Frish). 3. Akademiya nauk  
Belurusskoy SSR (for Yel'yashevich).  
(Spectrum analysis)

SOKOLOV, N.D.

35065

S/195/62/003/001/006/010  
E071/E136

5.1115

AUTHORS:

Kvlividze, V.I., Iyevskaya, N.M., Yegorova, T.S.,  
Kiselev, V.F., and Sokolov, N.D.

TITLE:

NMR studies of water vapour adsorption on the surface  
of silica gel

PERIODICAL: Kinetika i kataliz, v.3, no.1, 1962, 91-98

TEXT:

The mechanism of adsorption and the state of the adsorbed molecules on the surface of an adsorbent cannot be elucidated on the basis of purely adsorptive measurement. For this purpose some additional data on the system adsorbed substance - adsorbent obtained by physical methods are necessary. As a first stage in the investigations the authors studied signals of nuclear magnetic resonance from protons of hydroxyl groups of water adsorbed on the surface of silica gel. The results obtained were compared with adsorption properties of silica gel, with measurements of heats of adsorption and available spectroscopic data. Silica gels K-2 (K-2) and K-4 (K-4) obtained by the hydrolysis of  $\text{SiCl}_4$  and a purified sample of technical silica gel

Card 1/3

NMR studies of water vapour ...

S/195/62/003/001/006/010  
EC71/E136

KCK-3 (KSK-3) were used for the investigation. The NMR measurements were carried out at room temperature on a sample of 0.2-0.3 g. The width of the lines was measured as the distance between maxima on the differential curve. The second moment was calculated from the differential curve of the signal absorption. Additions of water vapour in the ampule with silica gel for NMR and adsorption measurements were carried out by means of a spring balance. Heats of adsorption were determined either directly from calorimetric measurements or by the differentiation of the curve relating the heat of wetting and the amount of water adsorbed on the specimen. From the adsorption data and heat of wetting curves, differential curves of the changes in free energy and entropy of adsorption were calculated. It was shown that molecules of water are absent on the surface of the samples evacuated at 200 °C. In the initial stage of adsorption a sharp decrease in the width of the line of the second moment was observed. These changes in the NMR signals agree with the trends of the curves of differential heat and entropy of adsorption. The possibility of interaction of water molecules

Card 2/3

NMR studies of water vapour ...

S/195/62/003/001/006/010  
EC71/E136

with the surface of silica gel through the coordination and hydrogen bonds is discussed.

There are 5 figures and 1 table.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im.  
M.V. Lomonosova, Fizicheskiy fakul'tet  
(Moscow State University imeni M.V. Lomonosov,  
Physics Division)

SUBMITTED: July 3, 1961

Card 3/3

X



S/030/62/000/010/006/007  
D204/D307

AUTHOR: Sokolov, N. D., Doctor of Physico-Mathematical Sciences

TITLE: Investigations of the hydrogen bond

PERIODICAL: Akademiya nauk SSSR. Vestnik, no. 10, 1962, 110-112

TEXT: A summary is given of a symposium on the subject of the hydrogen bond, held in Leningrad over July 4 - 7, 1962, organized by the Fizicheskiy fakultet Leningradskogo universiteta (Physics Faculty of Leningrad University), the Nauchnyy sovet (Scientific Council) concerned with the theories of chemical structure, kinetics, reactivity and catalysis, and by the Komissiya po spektroskopii Akademii nauk SSSR (Spectroscopy Commission of the Academy of Sciences, USSR), with the participation of over 200 scientists. The following topics were discussed: (1) The theory of the hydrogen bond; (2) Spectroscopy of the hydrogen bond (the most fully discussed subject during the symposium); (3) Structures of hydrogen-bonded compounds; (4) The nature of hydrogen bond in liquids

Card 1/2

MONOLOV, N.D., doktor fiz.-matem.nauk

Problems of quantum chemistry. Vest. AN SSSR 32 no.9:135-138 S  
'62. (MIRA 15:9)

(Quantum chemistry)

VETCHINKIN, S.I.; SOKOLOV, N.D.

Calculation of molecular integrals in quantum chemistry. Zhur.fiz.  
khim. 36 no.8:1754-1756 Ag '62. (MIRA 15:8)

1. Institut khimicheskoy fiziki AN SSSR i Moskovskiy gosudarstvennyy  
universitet fizicheskoy fakul'tet.  
(Quantum chemistry)

S/020/62/147/003/018/027  
B104/B186

AUTHORS: Korst, N. N., Savel'yev, V. A., Sokolov, N. D.

TITLE: Consideration of the averaging over the natural vibrational state when calculating the second moment of the nuclear magnetic resonance signal

PERIODICAL: Akademiya nauk SSSR. Doklady, v. 147, no. 3, 1962, 594 - 596

TEXT: Neglecting the uncertainty of the position of the protons in their natural vibrational state leads to the difference between the second moments of the n.m.r. proton line of ice as calculated with the van Vleck formula, which was  $\langle \Delta \omega^2 \rangle = 25 \text{ gauss}^2$  for stationary protons (this value is lower by 20%, when the protons perform a motion along the H-bonds) and those measured in experiments ( $T = 900\text{K}$ ;  $\langle \omega^2 \rangle = 36.7 \pm 1.7 \text{ gauss}^2$ ; K. Kume, J. Phys. Soc. Japan, 15, 1493 (1960)). The uncertainty is associated with the zero energy of the atoms. With consideration of neighboring proton pairs, the second moment of an ice single crystal is

$$\langle \Delta \omega^2 \rangle = \frac{3}{4} g^4 \beta^4 \hbar^{-2} I(I+1) \left[ \frac{3 \cos^2 \theta - 1}{R^3} \right]^2. \quad (2).$$

Card 1/3

Consideration of the...

S/020/62/147/003/018/027  
B104/B186

Expanding into powers of the deviations of the coordinates from their equilibrium values (after averaging over all possible mutual positions of the vectors  $\vec{R}$  and  $\vec{H}$ ) leads to

$$\langle \Delta \omega^2 \rangle = \frac{3}{5} g^4 \beta^4 \hbar^{-2} R^{-6} \{ (I+1) \left[ 1 + \frac{6 \overline{(\Delta r_1)^2}}{[2r_0 \sin \frac{\Phi_0}{2}]^2} + \frac{6 \overline{(\Delta r_2)^2}}{[2r_0 \sin \frac{\Phi_0}{2}]^2} + \frac{10 \overline{\Delta r_1 \Delta r_2}}{[2r_0 \sin \frac{\Phi_0}{2}]^2} + \frac{9}{4} \overline{(\Delta \Phi)^2} + \frac{87}{32} \overline{(\Delta \Phi)^2}^2 \right] \right\}, \quad (3)$$

for a polycrystal.  $\vec{R}_{ij}$  is the radius vector linking the  $i$ -th proton to the  $j$ -th. Computation with formula (3) showed that the term  $\overline{(\Delta \Phi)^2}$  makes up 70% of the correction to the second moment. Considering more distant protons one obtains

$$\langle \Delta \omega^2 \rangle = 0,80 \gamma^4 \hbar^2 R^{-6} \{ 1 + 0,21 \} = 31,1 \text{ gauss}^2.$$

Further improvement can be achieved by taking the hydrogen bond into  
Card 2/3

Consideration of the...

S/020/62/147/003/018/027  
B104/B186

account. The results show that proton transitions along the hydrogen bonds at  $T = 900^\circ\text{K}$  have little probability. Similar calculations of the second moments of polyethylene and of 1,2-dichloro ethane with formula (5) yielded good agreement with experimental data as published in J. Polym. Sci., 26, 171 (1957); J. Chem. Soc., 17, 972 (1949). There is 1 table.

ASSOCIATION: Fizicheskiy institut im. P. N. Lebedeva Akademii nauk SSSR  
(Physics Institute imeni P. N. Lebedev of the Academy of  
Sciences USSR)

PRESENTED: May 14, 1962, by V. N. Kondrat'yev, Academician

SUBMITTED: April 30, 1962

Card 3/3

SAVEL'YEV, V.A.; SOKOLOV, N.D.

Calculating the extension of the hydrogen bond in isotopic  
substitution in ice. Fiz. tver. tela 5 no.11:3273-3275 N  
'63. (MIRA 16:12)

1. Institut khimicheskoy fiziki AN SSSR, Moskva.

SZOKOLOV, N.D. [Sokolov, N.D.]

Quantum theory of the electron cloud of molecules. Magy fiz  
folyoir 11 no.2:159-176 '63.



SOKOLOV, N.D.

Symposium on hydrogen bonding. Opt. i spektr. 15 no.1:142-  
144 J1 '63. (MIRA 16:8)

(Hydrogen bonding)

SOKOLOV, N.D.

State and several problems of the theory of the electronic  
clouds of molecules. Usp.khim. 32 no.8:967-983 Ag '63.  
(MIRA 16:9)

1. Institut khimicheskoy fiziki AN SSSR.

SOKOLOV, N. D.

"Electron paramagnetic resonance" by S. A. Al'tshuler, B. M.  
Kozyrev. Reviewed by N. D. Sokolov. Usp. fiz. nauk 79 no.1:  
165-166 Ja '63. (MIRA 16:1)

(Paramagnetic resonance and relaxation)  
(Al'tshuler, S. A.) (Kozyrev, B. M.)

SOKOLOV, N. D.; KORST, N. N.; and SAVELYEV, B. A.

"Calculation of the Temperature-Dependence of Low Vibrational Frequencies in Ice."

papers presented by USSR and Polish Scientists at the VIIth European Congress on Molecular Spectroscopy (IUPAC), held 22-27 July 1963 in Budapest, Hungary.

SOKOLOV, N.D., prof., otv. red.; CHULANOVSKIY, V.M., prof., otv.  
red.; BUCHACHENKO, A.L., red.

[Hydrogen bonding] Vodorodnaia sviaz'; sbornik statei.  
Moskva, Izd-vo "Nauka," 1964. 339 p. (MIRA 17:8)

1. Akademiya nauk SSSR. Institut khimicheskoy fiziki.

ACCESSION NR: AP/028464

S/0181/64/006/004/1242/1243

AUTHORS: Korst, N. N.; Savel'yev, V. A.; Sekolov, N. D.

TITLE: The second moment of the NMR signal and the structure of ice

SOURCE: Fizika tverdogo tela, v. 6, no. 4, 1964, 1242-1243

TOPIC TAGS: nuclear magnetic resonance, second moment, ice structure

ABSTRACT: Magnitudes of the second moment  $\langle \Delta \omega^2 \rangle$  of the NMR signal from the protons of ice are computed by the modified Van Vleck formula in which the uncertainty in proton position due to the zero vibrational energy is considered. The extreme values of the geometrical parameters are used: the length of the O-H bond in ice,  $r_0$  (0.98 Å and 1.01 Å) and the H-O-H valence angle,  $\phi_0$  (104° and 109°28'). Values for the second moment are obtained between 33 and 40 (gauss)<sup>2</sup>, taking into account the frequency-temperature dependence and the exchange of protons by the tunnel effect. These values are not significantly different from the experimental value

$$\langle \Delta \omega^2 \rangle = 36.7 \pm 1.7 \text{ (gauss)}^2.$$

Card 1/2

ACCESSION NR: AP4028464

It is concluded that it is impossible to compute an exclusive model of ice in which there is an exchange of protons by random tunnelling. Orig. art. has: 6 equations.

ASSOCIATION: Institut khimicheskoy fiziki AN SSSR, Moscow (Institute of Physical Chemistry AN SSSR)

SUBMITTED: 25Nov63

DATE ACQ: 27Apr64

ENCL: 00

SUB CODE: PH

NO REF SOV: 004

OTHER: 005

Card 2/2

SAVEL'YEV, V.A.; SKOLOV, N.D.

Temperature dependence of the intermolecular vibration frequency in  
ice. Opt. i spektr. 17 no.1:35-37 J1 '64. (MIRA 17:9)



L 8758-65 ENT(1) IJP(1) ESD(1)/AEWL

ACCESSION NR: AP4044845

S/0051/64/017/003/0343/0348

AUTHOR: Pshenichnov, Ye. A.; Sokolov, N. D. 15

TITLE: Eigenvalues and probabilities of quantum transitions in a double asymmetrical potential well 16

SOURCE: Optika i spektroskopiya, v. 17, no. 3, 1964, 343-348

TOPIC TAGS: deuterated compound, quantum transition, proton transition, tunneling transition

ABSTRACT: The calculations were made for the A--D ... B bond, in analogy with the similar calculations made by R. L. Somorjai and D. F. Hornig (J. Chem. Phys., v. 36, 1980, 1962) for the A--H ... B bond. In addition to serving as a check on the characteristic features of a double asymmetrical well, the present calculations were aimed at improving the accuracy of the earlier ones, inasmuch as Somorjai and Hornig confined themselves only to the 20th order

Card 1/3

L 8758-65

ACCESSION NR: AP4044845

( $N = 20$ ) of the corresponding secular equation, which is not accurate enough. The accuracy is all the more suspect because the calculations of Somorjai and Hornig also lead to the conclusion that the tunneling transition of the proton from one well to the other is more probable at ordinary temperatures than above-the-barrier transition. The authors have calculated the eigenvalues and the eigenfunctions for  $N = 30$  for 4 variants of potential wells used by Somorjai and Hornig, using the electronic computer of the Mathematics Center of IKhF AN SSSR. The somewhat lesser attenuation of the oscillation intensity following substitution of deuterium for hydrogen, compared with the usual attenuation in free molecules, is used to conclude that a second potential well exists. It is noted in the mechanism of the transition of a proton or a deuteron from one potential well to the other, and that this investigation will be reported by the authors in the journal "Kinetika i kataliz." Orig. art. has: 3 tables, 1 figure, and 2 formulas.

Card 2/3

L 8758-65

ACCESSION NR: AP4044845

ASSOCIATION: None

SUBMITTED: 09Dec63

ENCL: 00

SUB CODE: GP

NR REF SOV: 002

OTHER: 008

Card 3/3

SOKOLOV, N.D.

Relation between the frequency shifts of valence and deformation vibrations in the hydrogen bridge. Opt. i spektr. 17 no.3:455-456 S '64.  
(MIRA 17:10)

SOKOLOV, N.D., doktor fiz.-matem.nauk

Conference on <sup>uc</sup>Advances in Quantum Chemistry held at Kishinev.  
Vest. AN SSSR 34 no. 1:108-111 Ja '64. (MIRA 17:5)

ALEKSEYEV, Igor' Vasil'evich; ALEKSEYEV, N.I., ed.

[Theory of nuclear magnetic resonance] Teoriya yadernogo  
magnitnogo rezonansa. Moskva, Nauka, 1964. 206 p.  
(BIA 17/10)

PSHENICHNIKOV, Ye.A.; SOKOLOV, N.D.

Mechanism of reactions involving proton transfer in polar  
solvents. *Kin. i kat.* 6 no.5:802-808 S=0 1965. (1965 18:11)

1. Institut khimicheskoy fiziki AN SSSR.

30 Kolov, N.F.

21 621.3.014.31 : 621.316.3  
 ✓ 2538. EXTINCTION OF A D.C. ARC IN A ROTATING MAGNETIC  
 FIELD A.E. Kaplyanskii, L.S. Polotovskii, N.F. Sokolov and  
 P.I. Petrov.

Elektrichestvo, 1956, No. 12, 29-32. In Russian.

An experimental investigation showed that the extinction of the arc by a rotating magnetic field of commercial frequency is in many cases more efficient, in others not less efficient, than arc-quenching by means of a constant or pulsating field. The supply of arc-quenching devices with 3-phase current does not offer difficulties, since in practice the direct-current concerned will almost exclusively be rectified 3-phase current. In some cases the use of a rotating field will reduce the dimensions of the arc-quenching device, this rendering the method suitable for use in d.c. switchgear, electronic converters and contact rectifiers. The latter use was tried out on a model of a mechanical 3-phase rectifier the contacts of which were placed in a rotating magnetic field. Without application of the field strong sparking at the contacts occurred at currents of only 0.5 A and 30 V. With the field applied, visible sparking did not occur up to the limiting current loading of 150 A of the model. The flux density of the rotating field must be 360-500 Gauss.

B.F. Kravtsov



SOKOLOV, N.F.; PINSKER, A.L.

Problems in the improvement of technical means for producing  
new tractors and agricultural machines. Trakt. i sel'khoz mash.  
no.12:28-34 D '58. (MIRA 11:12)

1. Nauchno-issledovatel'skiy institut Traktorosel'khoz mash.  
(Agricultural machinery industry)

SOKOLOV, N.F.

New automatic lines. Trakt.i sel'khoz mash. 30 no.2:38-39  
F '60. (MIRA 13:5)

1. Nauchno-issledovatel'skiy institut Traktorosel'khoz mash.  
(Agricultural machinery industry)

SOKOLOV, N.F.

Modernization of agricultural machinery plants. Trakt. i  
sel'khoz mash. 31 no.7:34-37 J1 '61. (MIRA 14:6)

1. Nauchno-issledovatel'skiy institut tekhnologii traktornogo i  
sel'skokhozyaystvennogo mashinostroyeniya.  
(Agricultural machinery industry)

LYSENKO, A.P.; SOKOLOV, N.F.; MIKHLIN, B.Z.

Methods of measuring weak magnetic fields in a wide spectrum  
of frequencies. Geofiz. prib. no.9:25-36 '61. (MIRA 15:11)  
(Magnetism, Terrestrial--Measurement)

SOKOLOV, N.G., inzhener.

Universal square tubular transverse beam. Rats. i izobr. predl. v  
stroi. no.130:13 '56. (MLRA 9:9)  
(Building blocks)

SOKOLOV, N.G., inzhener.

Device for filling vertical seams in large-block walls. Rats. 1  
izobr. predl. v stroi. no.130:14 '56. (MLRA 9:9)  
(Building blocks)

MAKSIMOVSKIY, N.P., kand.tekhn.nauk; SOKOLOV, N.G., inzh.

Making lightweight concretes for large panels and blocks using  
improved mixing machinery. Stroi. mat. 6 no.9:11-13 S '60.  
(MIRA 13:9)

(Mixing machinery) (Lightweight concrete)

GURICH, N. S., Eng.

Engr. Tech. Sci.

Dissertation: "Investigation of the Properties of a Series Motor Fed From a Series Transformer." Moscow Order of Lenin Power Engineering Inst named V. M. Molotov, 21 Nov 47.

SC: Vechernyaya Moskva, Nov, 1947 (Project #17836)



1899. Stability characteristics of a series motor fed from a series generator. SANDER, A. S. AND SEMULOV, M. G. *Elektricheskoe*, 10, 48-51 (Oct. 1946) 14. Russian. Series motors fed from series generators are frequently used for driving metal-cutting machines. Speed is regulated by varying the value of field diverter resistance; for speeds below that corresponding to full field on both machines, the generator field is shunted, for high speeds the motor field is weakened. Starting from Frolich's formula which gives the relation between flux and current an equation giving the relation between torque and speed of the motor is developed and illustrated by a graph. Curves giving motor r.p.m. against line current for various field to line current ratios ( $I_f/I_l$ ) have for speeds above the nominal and  $I_f/I_l > 1.5$  a max. to the left of which  $da/dI > 0$  and the circuit is unstable. The instability may occur in other cases in the nominal characteristic, and even when generator field is weakened. The instability is less frequent for identical machines than for dissimilar ones, the unstable conditions being more probable with saturated machines and those of higher output. The residual magnetism has an important influence upon the characteristics, its increase reducing the range of speed regulation.

SA B64C A.K.

ASAC-55A METALLURGICAL LITERATURE CLASSIFICATION

CONCLUSION

8(5)

AUTHORS:

Sirotin, Artemiy Afanas'yevich, Candidate SOV/161-58-2-24/30  
of Technical Sciences, Docent at Kafedra elektrooborudovaniya  
prompredpriyatiy Moskovskogo energeticheskogo instituta (Chair  
of Electrical Equipment of Industrial Enterprises), Sokolov,  
Nikolay Georgiyevich, Candidate of Technical Sciences, Docent  
at the Chair of Electrical Equipment of Industrial Enterprises,  
Moscow Power Engineering Institute, Rubtsov, Vladimir Vasil'yevich,  
Engineer at the 1 Podshipnikovyy zavod g. Moskvyy (1st Bearing  
Factory of the City of Moscow)

TITLE:

Electric Lag Drive of the Cross Feed (Transverse Feed) of  
Sphero-Grinders (Sledyashchiy elektroprivod poperechnoy podachi  
sferoshlifoval'nykh stankov)

PERIODICAL:

Nauchnyye doklady vysshey shkoly. Elektromekhanika i avtomatika,  
1958, Nr 2, pp 196 - 204 (USSR)

ABSTRACT:

The system of the cross feed electric drive should secure the  
removal of the main part of the supply at the maximum admissible  
actual feed. The slight rest of the supply has to be removed  
from a feed which secures the necessary quality of the product  
surface at minimum time. These requirements are met by the  
lag drive of the cross-feed which was developed for sphero-

Card 1/3

Electric Lag Drive of the Cross Feed (Transverse Feed)  
of Sphero-Grinders

SOV/161-58-2-24/30

grinders by the co-workers of the Moskovskiy energeticheskii institut (Moscow Power Engineering Institute) in cooperation with the Pervyy gosudarstvennyy podshipnikovyy zavod 1GP3 (First State Warehouse Factory 1GP3). The mode of effect of the lag system of cross feed is described and the electric wiring diagram of sphero-grinders with the lag drive of cross feed is shown. This system was fitted to the sphero-grinders of the Leningradskiy zavod imeni Il'icha (Leningrad Works imeni Il'ich) and to the machines of the Van-Norman works. The experimental investigation of the electric lag drive are described. The positive properties of the electric lag drive are as follows: 1) A check at the workshop has shown that this system meets the series production requirements of ball-bearing factories. 2) When correctly adjusted, the lag system prevents scrap of rings owing to burning. 3) The lag feed increases by efficiency a correct adjustment as compared to the existing mechanical facilities. 4) The surface quality at a lag feed is between the 7th and 8th class. 5) The lag feed permits an uncomplicated adjustment of one mode of operation to another. 6) The

Card 2/3

Electric Lag Drive of the Cross Feed (Transverse Feed) SOV/161-58-2-24/30  
of Sphero-Grinders

grinder is protected against excessive wear. 7) The electric diagram is not more complicated. 8) The diagram is more reliable than in other cases, due to the use of semiconductor valves instead of thermionic valves. 9) It is an automatic feed. 10) The specific energy consumption is lower by 16.1% as compared to mechanical feed. 11) The kinematic diagram of the cross feed assembly is by far less complicated. There are 8 figures.

ASSOCIATION: Kafedra elektrooborudovaniya prompredpriyatiy Moskovskogo energeticheskogo instituta (Chair of Electrical Equipment of Industrial Enterprises, Moscow Power Engineering Institute)

SUBMITTED: February 10, 1958

Card 3/3

SOKOLOV, Nikolay Georgiyevich; KLYUCHEV, V.I., kand. tekhn. nauk, retsenzent; KAPUNTSOV, Yu.D., inzh., retsenzent; ZIMIN, Ye.N., kand. tekhn. nauk, red.

[Design of electric drives for industrial mechanisms] Konstruirovaniye elektroprivodov proizvodstvennykh mekhanizmov; posobie dlia studentov spetsial'nosti [Elektrifikatsiia promyshlennykh predpriatii i ustanovok." Red.E.N.Zimin. Moskva, Mosk. energ.in-t, 1961. 222 p. (MIRA 16:6)  
(Electric driving)